

THE REPRESENTATIVITY OF PRETZEL KNOTS

MAKOTO OZAWA

In hope of recovery of my father

ABSTRACT. In the previous paper, the *representativity* of a knot K was defined as

$$r(K) = \max_{F \in \mathcal{F}} \min_{D \in \mathcal{D}_F} |\partial D \cap K|,$$

where $\mathcal{F}$ is the set of all closed surfaces of positive genus containing K and $\mathcal{D}_F$ is the set of all compressing disks for F in S^3 . Then an inequality $r(K) \leq b(K)$ has shown, where $b(K)$ denotes the bridge number of K . This shows that a (p, q, r) -pretzel knot has the representativity less than or equal to 3.

In the present paper, we will show that a (p, q, r) -pretzel knot has the representativity 3 if and only if (p, q, r) is either $\pm(-2, 3, 3)$ or $\pm(-2, 3, 5)$. We also show that a large algebraic knot has the representativity less than or equal to 3.

1. INTRODUCTION

Let K be a knot in the 3-sphere S^3 and F a closed surface of positive genus containing K . In [2], the *representativity* $r(F, K)$ of a pair (F, K) is defined as the minimal number of intersecting points of K and ∂D , where D ranges over all compressing disks for F in S^3 . Furthermore, we defined in [3] the *representativity* $r(K)$ of a knot K as the maximal number of $r(F, K)$ over all closed surfaces F of positive genus containing K . The representativity $r(K)$ of a knot K is a new knot invariant.

The following inequality is the first fundamental result for the representativity. Let $b(K)$ denote the bridge number of K .

Theorem 1.1 ([3, Theorem 1.2]). *For a knot K , $r(K) \leq b(K)$.*

This shows that a (p, q, r) -pretzel knot has the representativity less than or equal to 3. In the present paper, we characterize (p, q, r) -pretzel knots with the representativity 3.

Theorem 1.2. *Let K be a (p, q, r) -pretzel knot in S^3 . Then $r(K) = 3$ if and only if $(p, q, r) = \pm(-2, 3, 3)$ or $\pm(-2, 3, 5)$.*

Remark 1.3. It was shown in [1, Theorem III] that a (p, q, r) -pretzel knot is a torus knot if and only if $(p, q, r) = \pm(1, 1, 1)$, $\pm(-2, 3, 3)$ or $\pm(-2, 3, 5)$, and observed that a $\pm(-2, 3, 3)$ -pretzel knot is a $(3, \pm 4)$ -torus knot and a $\pm(-2, 3, 5)$ -pretzel knot is a $(3, \pm 5)$ -torus knot. In the proof of Theorem 1.2, we will give unknotted tori containing these pretzel knots directly.

2000 *Mathematics Subject Classification.* Primary 57M25; Secondary 57Q35.

Key words and phrases. pretzel knot, Montesinos knot, algebraic knot, representativity, bridge number.

Let K be a tangle composite knot, that is, which admits at least one essential tangle decomposition. We define the *tangle string number* $ts(K)$ of K as the minimal number of n over all n -string essential tangle decompositions of K .

Proposition 1.4 ([3, Example 3.6]). *For a tangle composite knot K , $r(K) \leq 2ts(K)$.*

By Proposition 1.4, $r(K) \leq 2$ if K is a composite knot, and $r(K) \leq 4$ if K has an essential Conway sphere. Nonetheless, the representativity of a large algebraic knot (i.e. algebraic knot with an essential Conway sphere) has more sharp upper bound.

Theorem 1.5. *Let K be a large algebraic knot. Then $r(K) \leq 3$.*

2. PROOFS

Proof. (of Theorem 1.2) Let K be a (p, q, r) -pretzel knot with a form of a Montesinos knot of type $(1/p, 1/q, 1/r)$. Thus K is obtained from three rational tangles (B_1, T_1) , (B_2, T_2) and (B_3, T_3) of slopes $1/p$, $1/q$ and $1/r$ respectively by connecting them in series. Let A be the “axis” of the Montesinos knot K , that is, an unknotted loop in the complement of K such that there exist three meridian disks D_1 , D_2 and D_3 in the solid torus $V = S^3 - \text{int}N(A)$ which cut (V, K) into three rational tangles (B_i, T_i) , (B_2, T_2) and (B_3, T_3) , where $B_1 \cap B_2 = D_1$, $B_2 \cap B_3 = D_2$ and $B_3 \cap B_1 = D_3$. See Figure 1.




FIGURE 1. The “axis” A of K and its exterior V

Suppose that $r(K) = 3$ and let F be a closed surface containing K which satisfies $r(F, K) = 3$. We assume that $|F \cap A|$ is minimal up to isotopy of F . Moreover, we assume that $|F \cap (D_1 \cup D_2 \cup D_3)|$ is minimal up to isotopy of F . Then, each component of $F \cap \partial V$ is a longitude of V which intersects ∂D_i in one point for each i , and $F \cap D_i$ consists of $|F \cap \partial V|/2$ -arcs and loops.

Claim 2.1. $|p|, |q|, |r| \geq 2$.

Proof. If $|p| \leq 1$, then K is a connected sum of two torus knots, a torus knot, or a 2-bridge knot. In either cases, we have $r(K) \leq 2$, a contradiction. $\square$

In the following, we show that $F \cap D_i$ has no loop and consists of mutually parallel $|F \cap \partial V|/2$ -arcs.

Claim 2.2. There exists no outermost arc α of $F \cap D_i$ with $|\alpha \cap K| = 0$.

Proof. Let α be an outermost arc of $F \cap D_i$ with $|\alpha \cap K| = 0$, and δ be the corresponding outermost disk. Since each component of $F \cap \partial V$ intersects ∂D_i in one point for each i , $\delta \cap \partial V$ connects two longitudinal components of $F \cap \partial V$. Hence by an isotopy of F along δ , $|F \cap A|$ can be reduced, a contradiction. $\square$

Claim 2.3. There exists no innermost loop β of $F \cap D_i$ with $|\beta \cap K| \leq 1$.

Proof. Let β be an innermost loop of $F \cap D_i$ with $|\beta \cap K| = 0$, and δ be the corresponding innermost disk. If $\partial\delta$ is essential in F , then $r(F, K) = 0$, a contradiction. Otherwise, $\partial\delta$ bounds a disk δ' in F and we can remove α from D_i by isotoping δ' into δ .

Next let β be an innermost loop of $F \cap D_i$ with $|\beta \cap K| = 1$, and δ be the corresponding innermost disk. Since $|\beta \cap K| = 1$, β is essential in F . Then $r(F, K) \leq 1$, a contradiction. $\square$

Claim 2.4. There exists no innermost loop β of $F \cap D_i$ with $|\beta \cap K| = 2$.

Proof. Suppose there exists an innermost loop β of $F \cap D_i$ with $|\beta \cap K| = 2$. Then β bounds a disk δ in F since $r(K) = 3$. Let β' be a loop of $\delta \cap (D_1 \cup D_2 \cup D_3)$ which is innermost in δ , and δ' be the corresponding innermost disk in δ . By Claim 2.3, $|\beta' \cap K| = 2$ and δ' contains one subarc of K . Since δ' is parallel into some D_i , at least one of p, q, r is equal to 0. This contradicts Claim 2.1. $\square$

By Claims 2.2, 2.3 and 2.4, $F \cap D_i$ has one of the following configurations. See Figure 2.

- (1) $F \cap D_i$ consists of mutually parallel arcs whose two outermost arcs α_{i1}, α_{i2} satisfying $|\alpha_{ij} \cap K| = 1$ for $j = 1, 2$.
- (2) $F \cap D_i$ consists of one arc α_i satisfying $|\alpha_i \cap K| = 2$.




FIGURE 2. Two configurations for $F \cap D_i$

In the following, we show that Configuration (2) does not exist.

Lemma 2.5. Let (B, T) be a trivial tangle and F is a properly embedded surface in B which contains T . Let $h : B \rightarrow \mathbb{R}_{\geq 0}$ be the standard Morse function with one critical point and suppose that each component of T has exactly one maximal point with respect to h . Then a pair (F, T) can be isotoped so that F has no inessential saddle point and each component of T has one maximal point with respect to h .

Proof. This lemma essentially follows [3, Lemma 2.1, 2.2]. $\square$

Claim 2.6. In Configuration (2), $F \cap B_1$ consists of a disk.

Proof. In Configuration (2), $F \cap \partial B_1$ consists of one loop β satisfying $|\beta \cap K| = 4$. If $F \cap B_1$ was not a disk, then by Lemma 2.5, there exists an essential saddle point of $F \cap B_1$ and this shows that $r(F \cap B_1, T_1) \leq 2$ as the proof of [3, Theorem 1.2]. $\square$

By Claim 2.6, we have $|p| \leq 1$. This contradicts Claim 2.1. Hence Configuration (2) does not exist. Hereafter, we assume Configuration (1). Since each component of $F \cap \partial V$ is a longitude of V which intersects ∂D_i in one point for each i , $F \cap \partial B_i$ has either following two cases. See Figure 3.

- (A) $F \cap \partial B_i$ consists of mutually parallel loops whose two innermost loops β_{i1}, β_{i2} satisfying $|\beta_{ij} \cap K| = 2$ for $j = 1, 2$.
- (B) $F \cap \partial B_i$ consists of one loop β_i satisfying $|\beta_i \cap K| = 4$.




FIGURE 3. Two cases for $F \cap \partial B_i$

Claim 2.7. In either cases, $F \cap B_i$ consists of disks.

Proof. In Case (A), without loss of generality, let δ be the innermost disk in ∂B_1 bounded by β_{11} . Since $r(F, K) = 3$, β_{11} bounds a disk δ' in F . Let γ be a loop of $\delta' \cap \partial B_1$ which is innermost in δ' , and ϵ be the corresponding innermost disk in δ' . Note that γ is a loop of $F \cap \partial B_1$, and ϵ contains exactly one subarc of K if and only if γ is either β_{11} or β_{12} .

If ϵ is contained in $B_2 \cup B_3$, then an algebraic tangle $(B_2 \cup B_3, T_2 \cup T_3)$ is inessential. It follows that $|q| \leq 1$ or $|r| \leq 1$ and it contradicts Claim 2.1. Hence ϵ is contained in B_1 .

If $\gamma = \beta_{11}$ and it bounds a disk ϵ in B_1 , then a loop of $F \cap \partial B_i$ which is next to β_{11} on ∂B_i , say β' , also bounds a disk ϵ' in B_1 such that $\epsilon' \cap F = \beta'$. Since $r(F, K) = 3$, β' bounds a disk ϵ'' in F and there is an isotopy sending ϵ'' to ϵ' . By the minimality of $|F \cap A|$ and $|F \cap D_i|$, β' bounds a disk ϵ' of $F \cap B_1$. Continuing this argument, each loop of $F \cap \partial B_1$ except for β_{12} bounds a disk of $F \cap B_1$ in B_1 . Now we apply Lemma 2.5 to (B_1, T_1) and $F \cap B_1$. If β_{12} does not bound a disk of $F \cap B_1$, then by Lemma 2.5, there exists an essential saddle point of $F \cap B_1$ and this shows that $r(K) \leq 1$, a contradiction. Hence each loop of $F \cap \partial B_1$ bounds a disk of $F \cap B_1$ if $\gamma = \beta_{11}$.

If $\gamma \neq \beta_{1j}$, then by the same argument as above, we can show that each loop of $F \cap \partial B_1$ bounds a disk of $F \cap B_1$.

In Case (B), we apply Lemma 2.5 to (B_i, T_i) and $F \cap B_i$. If $F \cap B_i$ was not a disk, then by Lemma 2.5, there exists an essential saddle point of $F \cap B_i$ and this shows that $r(F \cap B_1, T_1) \leq 2$, a contradiction. $\square$

We call $F \cap B_i$ *Type (A)* or *Type (B)* if $F \cap \partial B_i$ has Case (A) or Case (B) respectively. See Figure 4.

- (A) $F \cap B_i$ consists of mutually parallel k -disks whose two outermost disks E_{i1} and E_{i2} contain one string of T_i .
- (B) $F \cap B_i$ consists of one disk which contains two strings of T_i .




FIGURE 4. Two types for $F \cap B_i$

Claim 2.8. For Type (A), the boundary slope of $F \cap B_i$ is equal to the rational tangle slope of (B_i, T_i) .

Proof. This claim follows the definition of the slope of rational tangle. $\square$

Claim 2.9. For Type (B), let $1/m$ be the slope of a rational tangle and a/b be the boundary slope of $F \cap B_i$. Then one of the following conditions holds.

- (I) $am = b - 1$.
- (II) $a > 1$ and $am = b + 1$.


Proof. There are two possibility of the arrangement of T_i in $F \cap B_i$. Depending on them, we have two conditions (I) and (II). See Figure 5. We remark that it is required $a > 1$ in Condition (II). $\square$

Claim 2.10. In Claim 2.9, $a = 1$.

Proof. Since $F \cap B_i$ is a disk in B_i , for each region R between two consecutive arcs of $(F \cap B_i) \cap D_i$, there exists a disk δ in B_i such that $\delta \cap R$ is an arc connecting the two consecutive arcs and $\delta \cap (F \cap B_i) = \partial\delta - \text{int}(\delta \cap R)$. If $a \geq 2$, then $|\partial\delta \cap T_i| \leq 1$.

For the next tangle (B_j, T_j) which has a common disk D_i , we have a similar disk δ' in B_j as δ , for *some* region between two consecutive arcs of $(F \cap B_i) \cap D_i$, such that $|\delta \cap T_j| = 1$.

Then we have a disk $\Delta = \delta \cup \delta'$ such that $|\Delta \cap K| \leq 2$, a contradiction. $\square$



Condition (I) : slope 3/10 Condition (II) : slope 3/8 Condition (I) : slope 1/4

FIGURE 5. Two conditions for the boundary slope with the tangle slope 1/3

Hence, only (I) in Claim 2.9 occurs, and we have the following claim.

Claim 2.11. For Type (B), the boundary slope of $F \cap B_i$ is equal to $1/(m+1)$ if the slope of a rational tangle (B_i, T_i) is $1/m$.

Since $F \cap \partial V$ consists of longitudes, the next equation holds.

$$\frac{1}{p'} + \frac{1}{q'} + \frac{1}{r'} = 0, \quad (*)$$

where p' , q' and r' denote the boundary slopes of $F \cap B_1$, $F \cap B_2$ and $F \cap B_3$ respectively. Without loss of generality, we assume that $p' < 0$ and $q', r' > 0$. We note that the common denominator in the equation (*) is equal to $|F \cap D_i| = p'|F \cap B_1| = q'|F \cap B_2| = r'|F \cap B_3|$. In the following, we will give a condition for p', q', r' .

Lemma 2.12. *Generally, suppose that*

$$-\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 0 \quad (2 \leq a < b \leq c).$$

Then there exists a pair of coprime integers such that $k < l \leq 2k$ and an integer $d > 0$ such that

$$a = k(l-k)d, \quad b = l(l-k)d, \quad c = kld.$$

Proof. Put $a = km$, $b = lm$, where $k < l$ are coprime integers. Then

$$\begin{aligned} -\frac{1}{km} + \frac{1}{lm} + \frac{1}{c} &= 0 \\ -\frac{l-k}{klm} + \frac{1}{c} &= 0 \end{aligned}$$

Since $l - k$ divides m , putting $d = m/(l - k)$,

$$-\frac{1}{kld} + \frac{1}{c} = 0$$

Hence, $a = k(l - k)d$, $b = l(l - k)d$ and $c = kld$. Moreover, since $b \leq c$, we have $l \leq 2k$. $\square$

Claim 2.13. If $l = 2k$, then there exists a positive integer d such that $(p, q, r) = (p', q' - 1, r' - 1) = (-d, 2d - 1, 2d - 1)$. Here, $F \cap B_1$ is of Type (A) with $|F \cap B_1| = 2$, and $F \cap B_2$ and $F \cap B_3$ are of Type (B).

Proof. If $l = 2k$, then $l = 2$, $k = 1$ since l and k are coprime. Then we have $a = d$, $b = 2d$ and $c = 2d$ in Lemma 2.12. Since K is a knot, it follows that $q = 2d - 1$ and $r = 2d - 1$, thus $F \cap B_2$ and $F \cap B_3$ are of Type (B) by Claim 2.11. Then the common denominator in the equation (*) must be $2d$, $F \cap B_1$ is of Type (A) with $|F \cap B_1| = 2$. $\square$

Example 2.14. A $(-2, 3, 3)$ -pretzel knot is of this type in Claim 2.13.

Claim 2.15. If $l < 2k$, then there exists positive integers k, d such that $(p, q, r) = (p', q', r' - 1) = (-kd, (k + 1)d, k(k + 1)d - 1)$. Here, $F \cap B_1$ is of Type (A) with $|F \cap B_1| = k + 1$, $F \cap B_2$ is of Type (A) with $|F \cap B_2| = k$, and $F \cap B_3$ is of Type (B).

Proof. If $l - k$ is even, then both of a and b are also even. Since $b < c$, both of $F \cap B_1$ and $F \cap B_2$ are of Type (A) to have a common denominator in the equation (*). Thus by Claim 2.8, the slopes of rational tangles (B_1, T_1) and (B_2, T_2) coincide with those boundary slopes of $F \cap B_1$ and $F \cap B_2$. Therefore, p and q are even, this shows that K has more than two components.

Hence $l - k$ is odd, and one of l and k is even. It follows that one of a and b is even and c is even. Since $a < b < c$ and K has exactly one component, $F \cap B_3$ is of Type (B) by Claim 2.11. Then,

$$-\frac{1}{k(l - k)d} + \frac{1}{l(l - k)d} + \frac{1}{kld} = 0$$

$$-\frac{l - k}{kl(l - k)d} + \frac{1}{kld} = 0$$

Since $|F \cap D_2| = kl(l - k)d = kld$, $l - k = 1$ and $l = k + 1$. Hence $a = kd$, $b = (k + 1)d$ and $c = k(k + 1)d$, and $F \cap B_1$ is of Type (A) with $|F \cap B_1| = k + 1$, $F \cap B_2$ is of Type (A) with $|F \cap B_2| = k$ and $F \cap B_3$ is of Type (B). $\square$

Example 2.16. A $(-2, 3, 5)$ -pretzel knot is of this type in Claim 2.15, where $k = 2$ and $d = 1$.

Claim 2.17. In Claim 2.15, $d = 1$.

Proof. Suppose $d > 1$. There are $k(k + 1)d$ -arcs of $F \cap D_1$. Let α be the $k(k + 1)$ -th arc and α' be the $k(k + 1) + 1$ -th arc, and R be the region of D_1 between α and α' . Then there exists a disk δ in B_1 such that $\delta \cap (F \cap B_1)$ is an subarc β of $\partial\delta$ intersecting T_1 in one point and $\delta \cap R$ is an arc $\partial\delta - \text{int}\beta$ connecting α and α' . There also exists a disk δ' in B_2 such that $\delta' \cap (F \cap B_2)$ is an subarc β' of $\partial\delta'$ intersecting T_2 in one point and $\delta' \cap R$ is an arc $\partial\delta' - \text{int}\beta'$ connecting α and α' . Then a disk obtained from δ and δ' gives a compressing disk for F whose boundary intersects K in two points. This shows that $r(F, K) \leq 2$, a contradiction. $\square$

Therefore, we have two Types for the triple (p, q, r) .

- (1) $(p, q, r) = (-d, 2d - 1, 2d - 1)$ for some integer $d \geq 2$.
- (2) $(p, q, r) = (-k, (k + 1), k(k + 1) - 1)$ for some integer $k \geq 2$.

Claim 2.18. In Type (1), $d = 2$.

Proof. If $d \geq 3$, then we can find a compressing disk δ for F such that $|\partial D \cap K| = 2$, a contradiction. See Figure 6. $\square$




FIGURE 6. a compressing disk δ for F of Type (1) : $(-3, 5, 5)$ -pretzel knot

Claim 2.19. In Type (2), $k = 2$.

Proof. If $k \geq 3$, then we can find a compressing disk δ for F such that $|\partial D \cap K| = 2$, a contradiction. See Figure 7 for a $(-3, 4, 11)$ -pretzel knot. See Figure 8 for a $(-4, 5, 19)$ -pretzel knot. For $k \geq 5$, there exists a disk similar to Figure 8. $\square$

Conversely, let K be a $(-2, 3, 3)$ -pretzel knot. Then K has a closed surface F of Type (1), where $F \cap B_1$ is of Type (A) with $|F \cap B_1| = 2$ and $F \cap B_2$ and $F \cap B_3$ are of Type (B). See Figure 9. By the calculation of the Euler characteristic of F ,

$$\chi(F) = (2 + 1 + 1) - (4 + 4) + 4 = 0$$

we know that F is a torus, and there exist a meridian disks D and D' for each solid torus component obtained by splitting S^3 along F such that $|\partial D \cap K| = 3$ and $|\partial D' \cap K| = 4$. By checking that the algebraic intersection number of $\partial D \cap K$ and $\partial D' \cap K$ are equal to 3 and 4 respectively, it follows that K is a $(3, 4)$ -torus knot and F is an unknotted torus for K .

Next, let K be a $(-2, 3, 5)$ -pretzel knot. Then K has a closed surface F of Type (2), where $F \cap B_1$ is of Type (A) with $|F \cap B_1| = 3$, $F \cap B_2$ is of Type (A) with $|F \cap B_2| = 2$ and $F \cap B_3$ are of Type (B). See Figure 10. By the calculation of the Euler characteristic of F ,

$$\chi(F) = (3 + 2 + 1) - (6 + 6) + 6 = 0$$

we know that F is a torus, and there exist a meridian disks D and D' for each solid torus component obtained by splitting S^3 along F such that $|\partial D \cap K| = 3$ and $|\partial D' \cap K| = 5$. By checking that the algebraic intersection number of $\partial D \cap K$ and $\partial D' \cap K$ are equal to 3 and 5 respectively, it follows that K is a $(3, 5)$ -torus knot and F is an unknotted torus for K . $\square$




FIGURE 7. a compressing disk δ for F of Type (2) : $(-3, 4, 11)$ -pretzel knot

Proof. (of Theorem 1.5) Let K be a large algebraic knot. Then K admits an essential 2-string tangle decomposition $(B, T) \cup (B', T')$ by a Conway sphere S such that (B, T) is obtained from two rational tangles (B_1, T_1) and (B_2, T_2) by gluing along a disk D . Suppose that $r(K) = 4$ and let F be a closed surface containing K so that $r(F, K) = 4$.

We assume that $|F \cap S|$ is minimal up to isotopy of F . Then $F \cap S$ consists of one loop intersecting K in four points. Moreover, we assume that $|(F \cap B) \cap D|$ is minimal up to isotopy of $F \cap B$. Then $(F \cap B) \cap D$ consists of mutually parallel arcs whose two outermost arcs intersect T in one point (c.f. Configuration (1)) or one arc intersecting T in two points (c.f. Configuration (2)). These are similar configurations as in Figure 2.

In Configuration (1), let δ be the outermost disk in D bounded by an outermost arc of $(F \cap B) \cap D$. By cutting a disk in S bounded by $F \cap S$ along an arc $\delta \cap S$, and pasting one of the resultant subdisks and δ , we obtain a compressing disk δ' for F such that $|\partial\delta' \cap K| \leq 3$, a contradiction.

In Configuration (2), $F \cap \partial B_i$ consists of one loop. By applying Lemma 2.5 to $F \cap B_i$, there exists a compressing disk ϵ for $F \cap B_i$ such that $|\partial\epsilon \cap K| \leq 2$ or $F \cap B_i$ is a disk. In the former case, we have a contradiction. Hence $F \cap B_i$ is a disk and $F \cap B$ is also a disk. However, this shows that a tangle (B, T) is inessential, a contradiction. $\square$




FIGURE 8. a compressing disk δ for F of Type (2) : $(-4, 5, 19)$ -pretzel knot




FIGURE 9. a $(-2, 3, 3)$ -pretzel knot and an unknotted torus for a $(3, 4)$ -torus knot




FIGURE 10. a $(-2, 3, 5)$ -pretzel knot and an unknotted torus for a $(3, 5)$ -torus knot

REFERENCES

1. A. Kawauchi, *Classification of pretzel knots*, Kobe J. Math. **2** (1985) 11–22.
2. M. Ozawa, *Non-triviality of generalized alternating knots*, J. Knot Theory and its Ramifications **15** (2006) 351–360.
3. M. Ozawa, *Bridge position and the representativity of spatial graphs*, arXiv:0909.1162 (2009).

DEPARTMENT OF NATURAL SCIENCES, FACULTY OF ARTS AND SCIENCES, KOMAZAWA UNIVERSITY, 1-23-1 KOMAZAWA, SETAGAYA-KU, TOKYO, 154-8525, JAPAN

Current address: Department of Mathematics and Statistics, The University of Melbourne Parkville, Victoria 3010, Australia (temporary)

E-mail address: w3c@komazawa-u.ac.jp, ozawam@unimelb.edu.au (temporary)